



## **ANNUAL REPORT EXCERPT**

Ten Most Common Types of Invasive Cancer, Counts by Race and Ethnicity, 2013-2017

Breast cancer was the most common cancer diagnosed among females. Colon and rectum cancer was the third most common.

Lung and bronchus cancer was the second most common cancer among Black and White females, while thyroid cancer was the second most common cancer among Asian/Pacific Islander and Hispanic females.

Cervical cancer, a preventable cancer, was among the ten most common cancers for Black and Hispanic females.

Rank	Asian/ Pacific Islander	Black	White	Hispanic*
1	Breast	Breast	Breast	Breast
	(2,414)	(4,874)	(30,501)	(4,032)
2	Thyroid	Lung and Bronchus	Lung and Bronchus	Thyroid
	(537)	(1,713)	(13,360)	(1,111)
3	Colon and Rectum	Colon and Rectum	Colon and Rectum	Colon and Rectum
	(472)	(1,451)	(8,495)	(1,102)
4	Corpus and Uterus,	Corpus and Uterus,	Corpus and Uterus,	Corpus and Uterus,
	NOS	NOS	NOS	NOS
	(436)	(1,188)	(7,770)	(964)
5	Lung and Bronchus,	Thyroid	Thyroid	Lung and Bronchus
	(414)	(588)	(5,497)	(909)
6	Ovary (242)	Pancreas (538)	Melanoma of the Skin (4,581)	Non-Hodgkin Lymphoma (600)
7	Non-Hodgkin Lymphoma (205)	Non-Hodgkin Lymphoma (463)	Non-Hodgkin Lymphoma (4,469)	Cervix Uteri (428)
8	Stomach (169)	Myeloma (437)	Pancreas (3,230)	Kidney and Renal Pelvis (394)
9	Pancreas (140)	Kidney and Renal Pelvis (417)	Urinary Bladder** (2,812)	Leukemia (393)
10	Leukemia	Cervix Uteri	Leukemia	Pancreas
	(135)	(373)	(2,807)	(387)

## **New Jersey Females**

Source: New Jersey State Cancer Registry October 2019 submission file, New Jersey Department of Health. This table is color-coded by cancer type. Counts are inside parentheses.

\*Persons of Hispanic ethnicity may be of any race or combination of races. The categories of race and ethnicity are not mutually exclusive.

\*\*Includes in situ.

The New Jersey State Cancer Registry (NJSCR) Data Guidelines containing comprehensive data analysis methods, race and ethnicity classification, population and mortality data sources, and additional information related to NJSCR data can be found at <a href="https://www.nj.gov/health/ces/reports.shtml">https://www.nj.gov/health/ces/reports.shtml</a>.

Ten Most Common Types of Invasive Cancer, Counts by Race and Ethnicity, 2013-2017

## **New Jersey Males**

Rank Asian/ Black White Hispanic\* **Pacific Islander** Prostate Prostate Prostate Prostate 1 (1, 148)(5,373) (24, 924)(3.348)**Colon and Rectum** Colon and Rectum Lung and Bronchus Lung and Bronchus 2 (1,204)(585) (1,572) (11,993)3 Lung and Bronchus **Colon and Rectum Colon and Rectum** Lung and Bronchus (540) (1,349)(8,993) (939) Non-Hodgkin **Kidney and Renal Urinary Bladder\*\*** Non-Hodgkin 4 Lymphoma Pelvis (8, 155)Lymphoma (258) (656) (713) **Oral Cavity and Urinary Bladder\*\*** Melanoma of the **Kidney and Renal** 5 Pharynx (503) Skin Pelvis (255) (6,458) (613) **Urinary Bladder\*\* Urinary Bladder\*\*** 6 Non-Hodgkin Non-Hodgkin (233)Lymphoma Lymphoma (596) (462) (5,200)7 Stomach **Kidney and Renal** Leukemia Pancreas (228)(437) Pelvis (492) (4, 548)Liver Liver Leukemia Liver 8 (216)(435)(3,747)(399)**Oral Cavity and** 9 **Kidney and Renal** Myeloma Stomach Pelvis (391) (391) Pharynx (200) (3,502)10 Leukemia **Oral Cavity and** Pancreas **Oral Cavity and** (187) Pharynx (3,220) Pharynx (375) (350)

Source: New Jersey State Cancer Registry October 2019 submission file, New Jersey Department of Health. This table is color-coded by cancer type. Counts are inside parentheses.

\*Persons of Hispanic ethnicity may be of any race or combination of races. The categories of race and ethnicity are not mutually exclusive.

\*\*Includes in situ.

Prostate cancer was the most common cancer diagnosed among New Jersey males.

Colon and rectum cancer was the second most common cancer for Asian/ Pacific Islander and Hispanic males, and the third most common for Black and White males.

Lung and bronchus cancer was the second most common cancer among Black and White males and the third most common cancer among Asian/Pacific Islander and Hispanic males.